



Physical Chemistry Chemical Physics

# Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: R. Metzler and M. Weiss, *Phys. Chem. Chem. Phys.*, 2025, DOI: 10.1039/D5CP01378J.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the Information for Authors.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.





Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM

# Discriminating stochastic processes for the assessment of materials<sup>Online</sup> properties by diffusion measurements

Ralf Metzler

Institute of Physics & Astronomy, University of Potsdam, Karl-Liebknecht-St 24/25, 14476 Potsdam, Germany and Asia Pacific Centre for Theoretical Physics, Pohang, 37673, Republic of Korea

Matthias Weiss\*

Experimental Physics I, University of Bayreuth, Universitätsstr. 30, D-95447 Bayreuth, Germany

# Abstract

Assessing the materials properties of complex media, e.g. colloidal suspensions or intracellular fluids, frequently relies on quantifying the diffusive motion of tracer particles. In particular, from the particles' mean squared displacement (MSD) one may infer the complex shear modulus of the medium. Yet, experimentally the same power-law forms of the MSDs emerge for tracer diffusion in very different environments. For example, diffusive motion in a static maze of fractal obstacles (obstructed diffusion, OD) and motion in viscoelastic fluids (often described by fractional Brownian motion, FBM) can show an identical sublinear MSD scaling, but an MSD-derived complex shear modulus is meaningless for OD as the system does not feature any viscoelasticity. Here we show that OD and FBM trajectories are highly similar in many observables, including the MSD and the autocovariance function that reports on the memory of the particle motion. The Gaussianity and/or the asphericity of trajectories, extracted with single-particle tracking, allows for a proper discrimination of OD and FBM, facilitating a meaningful interpretation of the materials properties of the medium. In contrast, techniques that only monitor particle number fluctuations in a region of interest are not capable of discriminating highly similar random processes like FBM and OD as they only rely on the MSD. We therefore highly recommend the use of the more informative tracking of single particles when aiming to asses materials properties of the medium under investigation.

<sup>\*</sup>Corresponding author: matthias.weiss@uni-bayreuth.de

#### I. INTRODUCTION

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM.

View Article Online DOI: 10.1039/D5CP01378J

Quantifying diffusive motion is a versatile and often indispensible tool for determining the (heterogeneous) materials properties of complex fluids, from artificial systems such as liquid crystals, colloidal suspensions, and (semi)dilute polymer solutions to fluids in living cells, e.g., the cytoplasm and nucleoplasm. In the most basic scenario, evaluating the Brownian motion of spherical tracer particles with radius R in an isotropic fluid at thermal equilibrium yields a diffusion constant D, from which the local viscosity  $\eta$  can be retrieved via the Stokes-Einstein relation  $D = k_B T/(6\pi\eta R)$  [1]. Recent all-atom molecular dynamics simulations demonstrate that a time-local analog of the Stokes-Einstein relation even exists for simple proteins with fluctuating shapes in water [2].

Considering more complex samples, e.g. viscoelastic fluids, the diffusive transport often features anomalous characteristics in the sense that the mean-squared displacement (MSD) of tracer particles does not increase linearly in time but frequently shows the sublinear, power-law scaling  $\langle \mathbf{r}^2(\tau) \rangle \propto K\tau^{\alpha}$  with  $\alpha < 1$ , a phenomenon called "subdiffusion" [3]. Several stochastic processes can yield subdiffusive motion, with some processes even showing signatures of weak ergodicity breaking (see [4] for review). For conciseness we will restrict ourselves here to processes with stationary increment statistics that can be linked directly to materials properties at thermal equilibrium. The generalized diffusion coefficient K has units of area per fractional time and only becomes identical to the familiar diffusion constant D for  $\alpha = 1$  ("normal diffusion"). The sublinear MSD scaling and the unconventional units of K reflect the multi-scale nature of the fluid's materials properties that go beyond a simple constant viscosity. In such cases, the complex shear modulus  $G(\omega) = G'(\omega) + iG''(\omega)$  is an informative and more extensive measure that reports on the fluid's elastic (G') and viscous (G'') response when shearing it at frequency  $\omega$  [5].

At thermal equilibrium, the MSD of tracer particles may actually be used to determine the complex shear modulus  $G(\omega)$  via a Laplace transformation and an analytical continuation [6]. It is worth noting, however, that this approach tacitly assumes that the particles' random motion is indeed governed by the fluid's viscoelasticity, hence causing a non-trivial MSD due to the viscoelastic material property. Supposedly the best known stochastic model for describing diffusion in viscoelastic environments is fractional Brownian motion (FBM) in its subdiffusive form [7]. FBM is a non-Markovian Gaussian stochastic process with an anti-

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM

persisent memory that is set by a single parameter, the Hurst coefficient  $H \underset{\text{O} \in H^{\text{ievArt}}(d) \to 0}{\text{O} \otimes (10,103)/\text{DSCPOI378J}}$  for subdiffusion). The Hurst coefficient determines the MSD scaling exponent as  $\alpha = 2H$ . Translating this MSD into the complex shear modulus yields the non-trivial scaling  $|G(\omega)| \propto G'(\omega) \propto G''(\omega) \propto \omega^{\alpha}$  reporting on the viscous and elastic material properties.

However, experimentally acquired trajectories of tracers in a yet uncharacterized fluid may show a sublinear MSD scaling that is not related to viscoelasticity at all, hence jeopardizing a meaningful interpretation in terms of the MSD-derived complex shear modulus. This may already occur for Brownian diffusion, when the ("static") noise from inaccuracies in determining the particle position effect an apparent subdiffusion at shorter times [8, 9]. Another prime example is "obstructed diffusion" (OD), when particles move in a fractal maze of (immobile) obstacles: Randomly placing immobile obstacles with a density near or at the percolation threshold is known to result in a long-lasting or even asymptotically long subdiffusion of tracers as they can only explore a fractal subset of space [10, 11]. Although this scenario creates a stationary and subdiffusive random motion, the system contains no viscoelastic medium at all. Therefore, calculating  $G(\omega)$  from the MSD would erroneously suggest that the particles moved in a viscoelastic fluid, albeit this was not the physical nature of the observed random motion. This situation becomes even more critical in cases when a macroscopic rheological assessment, i.e., an alternative means to assess  $G(\omega)$ , cannot be used to complement the diffusion measurements—as the two approaches probe different length scales. An example is the motion of tracers in a fully polymerized hydrogel that features a typical mesh size, that is similar to the diameter of the tracer particle. While here the tracer may report a sublinear scaling of the MSD due to OD, macroscopic rheology will only report a rubber-like elasticity without any viscosity. It is therefore expedient to extract all necessary information from the diffusion measurement itself when determining the materials properties of a complex fluid—yet without falling into the trap of misinterpreting the data by focusing solely on the MSD.

Here we demonstrate that this may a priori be somewhat delicate, even when dealing with spatiotemporally homogenous systems. Namely, we show that FBM and OD display very similar features in surprisingly many experimentally accessible observables, e.g., a sublinear MSD and a distinct anti-persistent autocovariance function, hence impeding a simple discrimination of the two scenarios. We find, however, that a detailed analysis of an ensemble of trajectories can be used to identify FBM, hence supporting a proper interpretation of an

Physical Chemistry Chemical Physics Accepted Manuscrip

MSD-derived complex shear modulus via single-particle tracking experiments. In contraist<sup>Online</sup> Doi:10.1039/D5CP01378J ensemble-based measurement techniques that do not yield individual trajectories basically only exploit the MSD and are hence inadequate for distinguishing scenarios that have very similar properties like FBM and OD. We emphasize this aspect by considering techniques that rely on fluctuating particle numbers in a fixed observation volume. From our data we advocate single-particle tracking as the method of choice if one wishes to arrive at a meaningful interpretation when translating MSDs to complex shear moduli.

# II. METHODS

To simulate OD random walks, we used two-dimensional random mazes on a square lattice with an obstacle density of  $\phi = 40\%$ , close to the percolation threshold of obstacle percolation [11]. Specifically, we randomly chose 40% inaccessible sites on a 400 × 400 square lattice with periodic boundary conditions in which ten tracers were allowed to move according to the blind ant algorithm (i.e., 100% probability for nearest-neighbor-hopping attempt, acceptance only if new site is not blocked by an obstacle). Tracers were treated as ghosts that do not see each other but only interact with the impenetrable obstacles. Moreover, tracers were checked to not leave the first unit cell of the lattice on average even in the most mobile case, i.e., finite size effects due to the periodic boundaries were negligible. In total, 100 runs of this setting were performed, yielding a total of M = 1000 trajectories. Each run consisted of  $5 \times 10^6$  sweeps (all particles try to move once per sweep), and positions were stored every 100th step, i.e., the trajectory length was fixed to  $N = 5 \times 10^4$ .

To also consider non-static mazes, the same amount of runs were performed with obstacles moving according to the blind ant algorithm every Qth sweep ( $Q = 10^3, 10^4, 10^5$ ). For a movement attempt of an obstacle, all other obstacles and the tracers were treated as impenetrable to avoid inconsistencies. For comparison, an ensemble of FBM trajectories with the same statistics and a Hurst coefficient H = 0.35 was obtained as described previously [12]. FBM trajectories of length N with a scrambled memory kernel were obtained by concatenating independent FBM trajectories with only 50 positions, i.e., every 50 time steps the memory is randomized for the next step increment. For the analysis of these ensembles of trajectories, we used our recently introduced toolbox of Matlab routines [12].

Time steps and lattice constants for OD were adjusted to reach experimentally reasonable

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM.

values. To this end, the time interval between successive positions in the analyzed VierArie Control to the second state of the transmission of t

# **III. RESULTS AND DISCUSSION**

Starting from the fact that FBM (with an appropriately chosen Hurst coefficient) and OD in a fractal percolation cluster have the same sublinear scaling of the MSD, we wanted to test by means of simulations in which (experimentally accessible) observables a clear discrimination between the two random motions is possible. For technical simplicity, we restricted our simulations to tracer motion in two dimensions when simulating FBM trajectories and OD tracer motion in random mazes with an obstacle density close to the percolation threshold (see Methods for details).

#### A. Mean-squared displacement

In agreement with the literature on diffusion in static fractal percolation clusters [10, 13], we observed a subdiffusive MSD  $\langle \mathbf{r}^2(\tau) \rangle_{t,E} \propto \tau^{\alpha}$  with  $\alpha \approx 0.7$  (Fig. 1) for OD in a maze of immobile obstacles. The same sublinear MSD scaling is obtained for FBM with a Hurst cofficient H = 0.35 [7]. Hence, if only an experimentally determined MSD is at hand, one cannot decide which of the two processes is underlying the data.

Upon mobilizing obstacles, i.e., when updating obstacle positions every Qth step, the subdiffusive scaling of the emerging OD was seen to become transient (Fig. 1): Beyond a crossover time  $t_c \propto Q$  the trivial scaling of normal diffusion ( $\alpha = 1$ ) was seen to emerge, in line with previous results [14]. On time scales below  $t_c$  the tracers therefore experienced an effectively immobile maze of obstacles, akin to a static percolation cluster, while on significantly longer time scales the tracers and obstacles were equally mobile, hence yielding  $( au)
angle_{\mathsf{t},\mathsf{E}} [\mu\mathsf{m}^2]$ 

10<sup>0</sup>

Physical Chemistry Chemical Physics Accepted Manuscrip



 $\tau$  [S]

10<sup>2</sup>

View Article Online DOI: 10.1039/D5CP01378J



10<sup>1</sup>

the limit of a hard-sphere gas with normal diffusion.

10<sup>3</sup>

10<sup>2</sup>

10

10<sup>0</sup>

10

10

Such a transient subdiffusion can also be obtained when using FBM trajectories with H = 0.35 but intermittently resetting or scrambling the memory kernel (cf. Methods). An example for the resulting transient subdiffusion, being highly similar to OD with  $Q = 10^3$ , is shown in Fig. 1. Therefore, OD and FBM also show highly similar MSDs when being forced to have only a transient subdiffusion characteristics. We note that a similar crossover to normal diffusion can be observed for "tempered FBM", in which the power-law correlations of the noise include exponential or steeper power-law cut-offs [15].

## B. Autocovariance function

Given that FBM is a non-Markovian stochastic process by definition, we next probed potential differences of FBM and OD in the (normalized) autocovariance function (ACVF),

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM.

#### **Physical Chemistry Chemical Physics**

also known as velocity autocorrelation function,

$$C(\xi) = \left\langle \frac{\langle \mathbf{v}(t)\mathbf{v}(t+\tau)\rangle_t}{\langle \mathbf{v}(t)^2 \rangle_t} \right\rangle_E.$$
 (1)

Here, the instantaneous velocity is defined via the spatial increment taken within a time interval  $\delta t$ ,  $\mathbf{v}(t) = (\mathbf{r}(t + \delta t) - \mathbf{r}(t))/\delta t$ , and the lag time  $\tau$  is rescaled as  $\xi = \tau/\delta t$ . As a result, we observed that the ACVF of OD in a static maze shows a pronounced anti-persistent dip at around  $\xi = 1$ , irrespective of the choice of  $\delta t$  (Fig. 2a). In fact, this signature of an anti-persistent memory matches a previous report [16]. More surprising, however, is the fact that the analytical FBM expression for the ACVF,

$$C_{\rm FBM}(\xi) = \frac{1}{2} \Big[ (\xi + 1)^{\alpha} + |\xi - 1|^{\alpha} - 2\xi^{\alpha} \Big],$$
(2)

with  $\alpha = 2H = 0.7$  fits these numerical data for OD so well that one would confuse it with an ensemble of FBM trajectories at H = 0.35 if oblivious to the different origin of the data. Experimental trajectories therefore would not allow one to discriminate OD in a static maze from FBM by means of the ACVF.

ACVFs for OD in a mobilized maze still agree mostly with the analytical FBM expression, but significant deviations become visible for  $\xi > 1$  (see Fig. 2a). The same holds true for FBM trajectories with a scrambled memory kernel. Therefore, the ACVF cannot be used to clearly discriminate OD and FBM, albeit one may have expected this. In Fig. 2b, the correlation decay from the pronounced minimum at  $\xi = 1$  towards zero is highlighted by the double-logarithmic axes. For FBM,  $|C(\xi)| \propto \xi^{\alpha-2}$  is expected. Again, OD in a static maze follows this power law with a remarkable accuracy, whereas OD with mobile obstacles yields deviations that increase with increasing obstacle mobility. Similar deviations from the power-law are also seen for FBM trajectories with a scrambled memory kernel, i.e., losses in the memory kernel for OD and FBM have similar effects on the ACVF in both cases.

The remarkably good agreement of Eq. (2) with data for OD in a static maze suggests that the ACVF possesses generic features that are the same for all random motions with stationary increment statistics. Indeed, it was already hypothesized earlier [16] that the ACVF decay  $|C(\xi)| \propto \xi^{\alpha-2}$  for  $\xi > 1$  is always observed for (anti-persistent) random motions that feature an MSD scaling  $\langle \mathbf{r}^2(\tau) \rangle_{t,E} \propto \tau^{\alpha}$ . An heuristic support of the hypothesis that the ACVF is just the second derivative of the MSD is given in App. A for a one-dimensional unbiased random walk process with stationary increments, such as OD or FBM, supplementing a previously discussed integral-based argument [15].

View Article Online

DOI: 10.1039/D5CP01378J



This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM.

FIG. 2: (a) The ACVF for OD in a static maze, as a function of the rescaled lag time  $\xi = \tau/\delta t$ , shows a pronounced anti-correlation (negative) dip and follows the same master curve for  $\delta t = 3, 5, 7\Delta t$ (open black squares, diamonds, circles). The data is, somewhat surprisingly, in very close agreement with the prediction Eq. (2) for FBM with  $H = \alpha/2 = 0.35$  (blue dash-dotted line). This suggests that OD in a static maze features a very similar scale-invariant anti-persistent memory as FBM. For OD in a mobile maze ( $Q = 10^3$ , red symbols; shifted upward for better visibility) the FBM prediction still yields a very good description albeit small deviations for  $\xi > 1$  become visible. FBM trajectories with Hurst coefficient H = 0.35 and a scrambled memory kernel follow the OD data for  $Q = 10^3$  (light-blue triangles). Therefore, the ACVF cannot be used to properly discriminate FBM and OD as both have highly similar properties. (b) For OD in a static maze (open black circles,  $\delta t = 7\Delta t$ ) the asymptotic power-law decay of the ACVF for  $\xi > 1$  also follows the FBM prediction  $|C(\xi)| \propto \xi^{\alpha-2}$  (blue dash-dotted line). For OD in a mobile maze, successively larger deviations are seen ( $Q = 10^3$ : red squares;  $Q = 10^4, 10^5$ : dark-red, grey lines). FBM trajectories with a scrambled memory kernel show similarly strong deviations (light-blue triangles) from the power law.

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM.

#### C. Power spectral density

View Article Online DOI: 10.1039/D5CP01378J

Going beyond the time domain, one might wonder whether the coefficient of variation for the trajectories' power spectral density (PSD) can be used to discriminate FBM from OD. The time- and ensemble-averaged PSD is known to follow the scaling  $S(f) \sim 1/f^{\alpha+1}$ for subdiffusive processes with stationary increment statistics [17]. For FBM, the PSD of individual trajectories also follows this scaling [17]. Fluctuations of single-trajectory PSDs around the ensemble-average are summarized by the coefficient of variation  $\gamma$ , the ratio of the standard deviation and the mean (see also [12]). While subdiffusive FBM has been shown to robustly yield  $\gamma = 1$  for virtually all frequencies f, normal diffusion generically yields  $\gamma = \sqrt{5}/2$  [17, 18]. Using the coefficient of variation, OD in a static maze can indeed be identified via its slow monotonic convergence to  $\gamma = 1$  for increasing frequencies (Fig. 3). However, OD in mobilized mazes assumes a behavior that is basically indistinguishable from FBM with a scrambled memory kernel: For large frequencies (corresponding to short times),  $\gamma \to 1$  is seen for OD in a maze of mobile obstacles, whereas for small frequencies (long times) a transient plateau with  $\gamma \to \sqrt{5}/2$  (the normal diffusion case) is observed. This is analogous to the change of the MSD scaling when translating the cross-over time to a frequency  $f_c = 1/t_c$ . Thus, for experimental purposes, also the coefficient of variation of the PSD is not suitable for a reliable discrimination of OD and FBM.

# D. Gaussianity and asphericity

Given that MSDs, ACVFs, and PSDs turn out to be inadequate quantities to discriminate subdiffusion in a viscoelastic medium (FBM) from subdiffusion in a fractal percolation maze (OD), one might wonder if there is any better quantity that is able to distinguish the two random motions. In the following, we will highlight deviations between the two processes in two observables, that are readily accessible in single-particle tracking experiments.

By construction, FBM is a Gaussian process whereas OD can be expected to show deviations from a Gaussian increment statistics [10, 11]. A versatile tool to quantify this aspect is the non-Gaussianity parameter (NGP) of the trajectories,

$$g(\tau) = \left\langle \frac{d}{3} \frac{\langle \mathbf{r}^4(\tau) \rangle_t}{[\langle \mathbf{r}^2(\tau) \rangle_t]^2} - 1 \right\rangle_E , \qquad (3)$$



View Article Online DOI: 10.1039/D5CP01378J

FIG. 3: The PSD coefficient of variation  $\gamma$  (shown as a function of the number of time steps, fT) strongly deviates from unity for OD in a static maze (black circles vs. blue dash-dotted line). In contrast, OD in a maze of mobile obstacles ( $Q = 10^3$ , red squares) shows an agreement with normal Brownian motion ( $\gamma = \sqrt{5}/2$ , black dashed line) for small frequencies (in line with  $\langle \mathbf{r}^2(\tau) \rangle \propto \tau$  for long lag times). The same is observed for FBM with a scrambled memory kernel (light-blue triangles), indicating that  $\gamma$  cannot reliably discriminate OD from FBM.

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM.

which tests whether the fourth and second moments of the increment statistic are independent of the time scale  $\tau$ . In Eq. (3), d refers to the spatial dimension (d = 2 in our case) and  $\langle . \rangle_E$  denotes an average over the ensemble of trajectories. By definition,  $g(\tau) = 0$  for a Gaussian process. As expected, the NGP for OD in a static maze shows strong deviations from zero, highlighting the clear non-Gaussian character of the random motion (Fig. 4a). Upon mobilizing the obstacles,  $g(\tau) \approx 0$  is regaind beyond the cross-over time  $t_c \propto Q$ , in line with the change observed in the MSD (cf. Fig. 1). Still, a clear non-zero NGP is seen below  $t_c$  for OD whereas FBM has a vanishing NGP on all times scales, even with a scrambled memory kernel (Fig. 3a). Thus, the Gaussianity of trajectories, which is easily available in single-particle tracking experiments, can indeed discriminate between FBM and OD.

We finally wondered whether the geometric shape of the acquired trajectories also can provide a robust means to distinguish FBM and OD, hence allowing for a meaningful interpretation of the apparent materials properties of the medium. For two-dimensional trajectories, the trajectory asphericity reads [19]

$$A = \frac{\langle (R_1^2 - R_2^2)^2 \rangle_E}{\langle (R_1^2 + R_2^2)^2 \rangle_E},\tag{4}$$

where  $R_1$  and  $R_2$  denote the gyration radii of individual trajectories that can be obtained

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM.



View Article Online DOI: 10.1039/D5CP01378J

FIG. 4: (a) The non-Gaussianity parameter (NGP) for OD in a static maze (black circles) strongly deviates from zero, highlighting that the random walk is not Gaussian on these time scales. For OD with mobile obstacles ( $Q = 10^3$ : red squares,  $Q = 10^4$ ,  $10^5$  dark-red and grey lines) a successive convergence to  $g(\tau) = 0$  is observed for  $\tau > t_c \propto Q$ , as expected already from the MSD scaling. FBM with a scrambled memory kernel shows a vanishing NGP throughout (light-blue triangles). (b) The asphericity of trajectory segments within a period  $\tau$  follows the expected  $A_{\text{FBM}} \approx 0.4$  for a pure FBM with Hurst coefficient H = 0.35 (blue dash-dotted line and asterisks) whereas normal Brownian motion features  $A_0 = 4/7$  [19] (indicated by black dash-dotted line). For OD (static: open black circles; mobile,  $Q = 10^3$ : red squares) the asphericity remains close to  $A_0$ , allowing one to discriminate OD from FBM. For FBM with a scrambled memory kernel (light-blue triangles) an interpolation from  $A_{\text{FBM}}$  to  $A_0$  for increasing trajectory length is observed, reflecting the crossover to a Markovian random walk for  $\tau \gg t_c$ . Concluding, OD and FBM can be discriminated by the values of A.

from the eigenvalues of the gyration tensor for each trajectory [12]. Straight rods and circle shapes yield A = 1 and A = 0, respectively, while trajectories of normal Brownian motion can be shown with mathematical rigor to feature  $A_0 = 4/7$  in two dimensions [19]. Subdiffusive FBM trajectories were shown to yield decreasing values of A for decreasing

Physical Chemistry Chemical Physics Accepted Manuscrip

Hurst coefficients, whereas OD in a static maze remains close to  $A_0$  [20]. It is therefore whereas  $A_0$  [20]. It is therefore with the sphericity is another experimentally accessible observable that can discriminate OD and FBM.

In line with this reasoning, we observed that OD in a static maze but also in the case of mobile obstacles, yielded trajectory asphericities that remained close to  $A_0 = 4/7$  (Fig. 3c). In contrast, a pure FBM with Hurst coefficient H = 0.35 yielded  $A \approx 0.4$ . For FBM with a scrambled memory kernel, an interpolation between this value and  $A_0$  is observed for increasing length of the trajectory segments (Fig. 4b), reflecting the crossover to a normal, Markovian random motion for  $\tau \gg t_c$ . In any case, OD and FBM can be discriminated via the asphericity of trajectories in a similarly robust fashion as seen for the NGP (cf. Fig. 4a). Both quantities are readily accessible with single-particle tracking methods. As will be shown in the next paragraph, ensemble-based techniques, that rely on monitoring the fluctuating number of particles in an observation volume, are not suited for revealing these subtle differences.

### E. Ensemble-based experimental approaches

There exist several ensemble-based techniques to assess the diffusive motion of tracers in a yet to be characterized medium. Techniques such as fluorescence recovery after photobleaching (FRAP) [21] only record the mean number of observable particles in a specified observation volume (region of interest, ROI), i.e., they monitor the relaxation of an observable back to its steady state after a perturbation. In FRAP experiments, this is done by quantifying the recovery of the mean fluorescence in an ROI after having bleached tracer particles in this region. More refined techniques do not require this invasive interaction with the sample but rely on monitoring fluctuating particle numbers in an ROI, an approach that was already utilized by Smoluchowski in his seminal work on diffusing colloids [22]. Monitoring particle number fluctuations instead of recording individual trajectories is, for example, a versatile approach when particles are too dense to be tracked properly or trajectories are too short to allow for a meaningful analysis along the lines described above.

Supposedly the most prominent and widespread technical implementation to exploit particle number fluctuations in a ROI is fluorescence correlation spectroscopy (FCS) [23]. Here, stationary fluctuations about a constant fluorescence,  $F(t) = \langle F \rangle + f(t)$ , are monitored with

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM.





FIG. 5: The squared particle number fluctuations  $\langle \Delta N(\tau)^2 \rangle_t$  for an OD in a static maze (black circles) and for a pure FBM with H = 0.35 (blue asterisks) fully overlap. The same is seen for OD in a maze of mobile obstacles ( $Q = 10^3$ : red squares) and FBM with a scrambled memory kernel (light-blue triangles). In both cases, the MSD-derived theoretical curve [Eq. (8)] fits the data (black line). Hence, FBM and OD cannot be discriminated.

high temporal resolution, allowing to extract the typical residence time  $\tau_D$  in the focus (i.e., in the ROI). The fluorescence autocorrelation function is then obtained via spatial averaging and reads (without normalization to the mean fluorescence)

$$C(\tau) = \langle F(t+\tau)F(t)\rangle_{t}$$
  
=  $\left\langle \int I(\mathbf{r})I(\mathbf{r}')\varrho(\mathbf{r},t+\tau)\varrho(\mathbf{r}',t)d(V,V')\right\rangle_{t}$   
=  $\int I(\mathbf{r})I(\mathbf{r}')G(\mathbf{r},\mathbf{r}',\tau)d(V,V'),$  (5)

where  $\rho(\mathbf{r}, t)$  denotes the particle density at position  $\mathbf{r}$  at time t and  $G(\mathbf{r}, \mathbf{r}', \tau) = \langle \rho(\mathbf{r}, t + \tau)\rho(\mathbf{r}', t)\rangle_t$  is the (diffusive) propagator of the particle density. For normal Brownian motion in one dimension,  $G(x, x', \tau) = \exp\left(-[x-x']^2/(4D\tau)\right)/\sqrt{4\pi D\tau}$ . In modern FCS approaches, particle counting is typically performed via the fluorescence signal from a confocal volume, approximating the relevant ROI  $I(\mathbf{r})$  as a Gaussian point-spread function. As a result, the fluorescence autocorrelation then decays algebraically with a typical time scale  $\tau_D$ , given by the effective area of the ROI divided by the diffusion constant [23]. Subdiffusion updates the exponent of the algebraic decay [24–26].

In the case that particles can be simply counted without the need to rely on their fluorescence signature, one can replace the ROI  $I(\mathbf{r})$  by a step function in every dimension,

Physical Chemistry Chemical Physics Accepted Manuscrip

yielding

View Article Online DOI: 10.1039/D5CP01378J

$$C(\tau) = \langle N(t+\tau)N(t)\rangle_t .$$
(6)

Since this is the only relevant autocorrelation function of the system, also the temporal variation of fluctuations of the squared particle number change is determined by this expression,

$$\langle \Delta N(\tau)^2 \rangle_t = \langle \{N(\tau) - N(t)\}^2 \rangle_t = 2 \langle N^2 \rangle_t - 2C(\tau) .$$
(7)

The latter quantity has recently been re-invented as "countoscope" and was used for the analysis of diffusive processes in dense colloidal systems [27]. For normal Brownian diffusion with diffusion constant D in d dimensions and a cubic box of edge length L, the analytical prediction is

$$\langle \Delta N^2(\tau) \rangle = 2 \langle N \rangle (1 - f(\tau)^d) \tag{8}$$

with

$$f(\tau) = \sqrt{\frac{4D\tau}{\pi L^2}} \left( e^{-L^2/(4D\tau)} - 1 \right) + \text{erf} \ , \left( \sqrt{L^2/(4D\tau)} \right).$$
(9)

where erf denotes the error function.

Given the structure of the formula and bearing in mind Smoluchowski's comments that the temporal evolution of  $\langle \Delta N(\tau)^2 \rangle$  is linked to the MSD, one may replace  $4D\tau$  by  $2\langle \mathbf{r}^2(\tau) \rangle / d$ in d dimensions, i.e., counting particles and inspecting the autocorrelator of the fluctuations is completely determined by the MSD. This is confirmed in Fig. 5, where FBM and OD data together with the MSD-determined theoretical curve is shown. Due to their very high similarity in their MSDs, FBM and OD cannot be discriminated—hence a proper assessment whether the medium has indeed a viscoelastic property remains obscure in such setups.

## IV. CONCLUSIONS

In summary, we conclude that monitoring fluctuations of particle numbers will only report on the MSD and hence will not allow one to retrieve the underlying stochastic process, even when dealing with spatiotemporally homogenous systems. Without this knowledge, however, the transfer from MSD to the complex shear modulus is prone to misinterpretations as, for example, OD is not associated with a viscoelastic medium (unlike FBM). In fact, the situation can become even more complicated: When the medium that is explored by the tracer cannot be approximated any more as a homogenous fluid (at thermal equilibrium) no

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM

Another central result in our study is the remarkable similarity between the ACVF of subdiffusive FBM and that of OD, in particular, the dip to negative (anti-persistent) values. If highly resolved data are available, there is the possibility to evaluate the area under the ACVF. For subdiffusive FBM, this should vanish identically to zero [15]. We also note that the ACVF shape, observed here for OD and subdiffusive FBM, also strongly resembles the ACVF of *confined* subdiffusive continuous time random walks [36], in which subdiffusion is effected by a scale-free probability density function (PDF) of immobilization times with an asymptotic power-law form  $\psi(\tau) \simeq \tau^{-1-\alpha}$  with  $0 < \alpha < 1$  [3]. Future studies may also benefit greatly from the statistics of mean-squared increments [37].

Our work was aimed at the evaluation of easily accessible observables. This may be complemented by other, more sophisticated data analysis, such as Bayesian methods [38, 39] or also deep learning-based approaches [40–44]. However, these are often not off-the-shelf solutions but rather require detailed knowledge on issues such as data pre-processing. Moreover, many of the available software suites do not contain all relevant stochastic processes, i.e. detailed tests like the one executed here on OD and FBM may require the implementation of such processes.

We finally note that the devlopment of FBM-type processes is still ongoing, even though FBM is by now more than 50 years old. First, there exist different definitions, including Mandelbrot's version in terms of a Weyl fractional integral [7], Lévy's definition via a Riemann-Liouville integral with initial non-stationarity [7, 45], and the Langevin equation

formulation with fractional Gaussian noise [7, 15]. While all three lead to the same by the politice politice of the parameters are chosen to vary, e.g., for a diffusing diffusivity [46]. In that case it can be shown that the associated PDF is also non-Gaussian for times below a typical correlation time. These phenomena will be analyzed in detail in the context of the present work in near future.

#### Acknowledgments

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM

MW gratefully acknowledges financial support by the VolkswagenStiftung (Az. 92738).

- Einstein A. Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen. Annalen der Physik. 1905;322:549-60.
- [2] Yamamoto E, Akimoto T, Mitsutake A, Metzler R. Universal Relation between Instantaneous Diffusivity and Radius of Gyration of Proteins in Aqueous Solution. Phys Rev Lett. 2021;126:128101.
- [3] Metzler R, Klafter J. The random walk's guide to anomalous diffusion: A fractional dynamics approach. Phys Rep. 2000;339:1-77.
- [4] Metzler R, Jeon JH, Cherstvy AG, Barkai E. Anomalous diffusion models and their properties: non-stationarity, non-ergodicity, and ageing at the centenary of single particle tracking. Phys Chem Chem Phys. 2014;16:24128-64.
- [5] Scott S, Weiss M, Selhuber-Unkel C, Barooji YF, Sabri A, Erler JT, et al. Extracting, quantifying, and comparing dynamical and biomechanical properties of living matter through single particle tracking. Phys Chem Chem Phys. 2023;25:1513-37.
- [6] Mason TG, Weitz DA. Optical Measurements of Frequency-Dependent Linear Viscoelastic Moduli of Complex Fluids. Phys Rev Lett. 1995;74:1250-3.
- [7] Mandelbrot BB, Ness JWV. Fractional Brownian Motions, Fractional Noises and Applications.
   SIAM Rev. 1968;10:422-37.
- [8] Meyer PG, Metzler R. Stochastic processes in a confining harmonic potential in the presence of static and dynamic measurement noise. New J Phys. 2023;25:063003.

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM

- [9] Martin MB D S Forstner, Käs JA. Apparent subdiffusion inherent to single particle tYackticle<sup>Online</sup> Biophys J. 2002;83:2109.
- [10] Havlin S, Ben-Avraham D. Diffusion in disordered media. Adv Phys. 1987;36:695-798.
- [11] Mardoukhi Y, Jeon JH, Metzler R. Geometry controlled anomalous diffusion in random fractal geometries: looking beyond the infinite cluster. Phys Chem Chem Phys. 2015;17:30134-47.
- [12] Rehfeldt F, Weiss M. The random walker's toolbox for analyzing single-particle tracking data. Soft Matter. 2023;19:5206.
- [13] Bouchaud JP, Georges A. Anomalous diffusion in disordered media: Statistical mechanisms, models and physical applications. Physics Reports. 1990;195(4-5):127-293.
- [14] Malchus N, Weiss M. Anomalous Diffusion Reports on the Interaction of Misfolded Proteins with the Quality Control Machinery in the Endoplasmic Reticulum. Biophys J. 2010;99:1321-8.
- [15] Molina-Garcia D, Sandev T, Safdari H, Pagnini G, Chechkin A, Metzler R. Crossover from anomalous to normal diffusion: truncated power-law noise correlations and applications to dynamics in lipid bilayers. New J Phys. 2018;20:103027.
- [16] Jacobs D, Nakanishi H. Autocorrelation functions for discrete random walks on disordered lattice. Phys Rev A. 1990;41:706-19.
- [17] Krapf D, Lukat N, Marinari E, Metzler R, Oshanin G, Selhuber-Unkel C, et al. Spectral Content of a Single Non-Brownian Trajectory. Phys Rev X. 2019;9:011019.
- [18] Sposini V, Krapf D, Marinari E, Sunyer R, Ritort F, Taheri F, et al. Towards a robust criterion of anomalous diffusion. Comm Phys. 2022;5:305.
- [19] Rudnick J, Gaspari G. The shapes of random walks. Science. 1987;237:384-9.
- [20] Ernst D, Hellmann M, Köhler J, Weiss M. Fractional Brownian Motion in Crowded Fluids. Soft Matt. 2012;8:4886-9.
- [21] Wachsmuth M. Molecular diffusion and binding analyzed with FRAP. Protoplasma. 2014;251:373.
- [22] von Smoluchowski M. Studien über Kolloidstatistik und den Mechanismus der Diffusion. Kolloid Z. 1916;18:48-54.
- [23] Rigler R. Fluorescence Correlation Spectroscopy. Springer Verlag (Berlin); 2001.
- [24] Weiss M, Hashimoto H, Nilsson T. Anomalous protein diffusion in living cells as seen by fluorescence correlation spectroscopy. Biophy J. 2003;84:4043-52.
- [25] Weiss M, Elsner M, Kartberg F, Nilsson T. Anomalous subdiffusion is a measure for cytoplas-

mic crowding in living cells. Biophys J. 2004;87:3518-24.

- [26] Szymanski J, Weiss M. Elucidating the Origin of Anomalous Diffusion in Crowded Fluids. Phys Rev Lett. 2009;103:038102.
- [27] Mackay EKR, Marbach S, Sprinkle B, Thorneywork AL. The Countoscope: Measuring Self and Collective Dynamics without Trajectories. Phys Rev X. 2024;14:041016.
- [28] Samanta N, Chakrabarti R. Tracer diffusion in a sea of polymers with binding zones: mobile vs. frozen traps. Soft Matt. 2016;12:8554-63.
- [29] Stadler L, Weiss M. Non-equilibrium forces drive the anomalous diffusion of telomeres in the nucleus of mammalian cells. New J Phys. 2017;19:113048.
- [30] Speckner K, Stadler L, Weiss M. Anomalous dynamics of the endoplasmic reticulum network. Phys Rev E. 2018;98:012406.
- [31] Kumar P, Theeyancheri L, Chaki S, Chakrabarti R. Transport of probe particles in a polymer network: effects of probe size, network rigidity and probe–polymer interaction. Soft Matt. 2019;15:8992-9002.
- [32] Sabri A, Xu X, Krapf D, Weiss M. Elucidating the Origin of Heterogeneous Anomalous Diffusion in the Cytoplasm of Mammalian Cells. Phys Rev Lett. 2020;125:058101.
- [33] Chubynsky MV, Slater GW. Diffusing Diffusivity: A Model for Anomalous, yet Brownian, Diffusion. Phys Rev Lett. 2014;113:098302.
- [34] Chechkin AV, Seno F, Metzler R, Sokolov IM. Brownian yet Non-Gaussian Diffusion: From Superstatistics to Subordination of Diffusing Diffusivities. Phys Rev X. 2017 Apr;7:021002.
- [35] Milster S, Koch F, Widder C, Schilling T, Dzubiella J. Tracer dynamics in polymer networks: Generalized Langevin description. J Chem Phys. 2024;160:094901.
- [36] Burov S, Metzler R, Barkai E. Aging and nonergodicity beyond the Khinchin theorem. Proc Natl Acad Sci USA. 2010;107:13228-33.
- [37] Wei Q, Wang W, Tang Y, Metzler R, Chechkin A. Fractional Langevin equation far from equilibrium: Riemann-Liouville fractional Brownian motion, spurious nonergodicity, and aging. Phys Rev E. 2025;111:014128.
- [38] Robson A, Burrage K, Leake MC. Inferring diffusion in single live cells at the single-molecule level. Phil Trans R Soc Lond B. 2013;368:20120029.
- [39] Thapa S, Lomholt MA, Krog J, Cherstvy AG, Metzler R. Bayesian analysis of single-particle tracking data using the nested-sampling algorithm: maximum-likelihood model selection ap-

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM

plied to stochastic-diffusivity data. Phys Chem Chem Phys. 2018;20:29018-37. View Article Online View Article Online 10.1039/D5CP01378J

- [40] Munoz-Gil G, Garcia-March MA, Manzo C, Martín-Guerrero JD, Lewenstein M. Single trajectory characterization via machine learning. New J Phys. 2020;22:013010.
- [41] Munoz-Gil G, Volpe G, Garcia-March MA, et al. Objective comparison of methods to decode anomalous diffusion. Nat Commun. 2021;12:6253.
- [42] Kowalek P, Loch-Olszewska H, Szwabiński J. Classification of diffusion modes in single-particle tracking data: Feature-based versus deep-learning approach. Phys Rev E. 2019;100:032410.
- [43] Seckler H, Metzler R. Bayesian deep learning for error estimation in the analysis of anomalous diffusion. Nat Commun. 2022;13:6717.
- [44] Seckler H, Szwabiński J, Metzler R. Machine-learning solutions for the analysis of singleparticle diffusion trajectories. J Phys Chem Lett. 2023;14:7910.
- [45] Lévy P. Random functions: general theory with special reference to Laplacian random functions. University of California Press, Berkeley; 1953.
- [46] Wei W, Chechkin AV, Metzler R. Different behaviors of diffusing diffusivity dynamics based on three different definitions of fractional Brownian motion; 2025. Available from: https://arxiv.org/abs/2504.19190.

View Article Online

DOI: 10.1039/D5CP01378J

#### Appendix A: Appendix

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM.

To obtain a heuristic argument, why an MSD scaling  $\langle \mathbf{r}^2(\tau) \rangle_{t,E} \propto \tau^{\alpha}$  may generically yield a ACVF decay  $|C(\xi)| \propto \xi^{\alpha-2}$  for  $\xi > 1$ , let us consider a one-dimensional unbiased random walk process with stationary increments and a time-averaged MSD with a power-law scaling

$$\langle x^2(\tau) \rangle_t = \langle [x(t+\tau) - x(t)]^2 \rangle_t = 2 \langle [x(t)]^2 \rangle_t - 2 \langle x(t+\tau)x(t) \rangle_t \propto \tau^{\alpha}$$

Mathematical random walk processes like the Wiener process are non-differentiable at every point, yet physical trajectories are continuous on (very) small time scales on which inertial effects and the impact of surrounding particles need to be treated with Newtonian mechanics. Only for sufficiently large time scales a simplified approximate description via an overdamped Langevin equation with uncorrelated noise becomes meaningful, yielding a non-differentiable random walk trajectory. We will therefore assume in the following that there is a small time intervall  $\Delta \tau$  for which the continuous Newtonian motion of the particle is still differentiable, so that derviatives can be approximated in a meaningful way by difference quotients. The second order derivative of the MSD hence reads

$$\frac{d^2 \langle \mathbf{r}^2(\tau) \rangle_t}{d\tau^2} = -2 \frac{d^2}{d\tau^2} \langle x(t+\tau)x(t) \rangle_t = -2 \left\langle \frac{d^2 x(t+\tau)}{d\tau^2} x(t) \right\rangle_t \\
= -2 \left\langle \frac{[x(t+\tau+\Delta\tau)+x(t+\tau-\Delta\tau)-2x(t+\tau)]x(t)}{\Delta\tau^2} \right\rangle_t \propto \tau^{\alpha-2} \quad (A1)$$

Using the definition of the ACVF [Eq. (1)] and abbreviating its (constant) normalization factor as  $v_0^2$ , one can relate  $C(\tau)$  to  $\langle \mathbf{r}^2(\tau) \rangle_t$ :

$$C(\tau) = \frac{\langle [x(t+\tau+\delta t) - x(t+\tau)][x(t+\delta t) - x(t)] \rangle_t}{v_0^2 \delta t^2}$$

$$= \frac{\langle x(t+\tau+\delta t)x(t+\delta t) \rangle_t + \langle x(t+\tau)x(t) \rangle_t - \langle x(t+\tau)x(t+\delta t) \rangle_t - \langle x(t+\tau+\delta t)x(t) \rangle_t}{v_0^2 \delta t^2}$$

$$= \frac{2\langle x(t+\tau)x(t) \rangle_t - \langle x(t+\tau-\delta t)x(t) \rangle_t - \langle x(t+\tau+\delta t)x(t) \rangle_t}{v_0^2 \delta t^2}$$

$$= \frac{\langle [2x(t+\tau) - x(t+\tau-\delta t) - x(t+\tau+\delta t)]x(t) \rangle_t}{v_0^2 \delta t^2}$$

where we have used again stationarity, i.e. invariance of averages with respect to shifts. Now assuming that  $\delta t \ll \tau$  (corresponding to  $\xi \gg 1$ ) the last line can be identified with Eq. (A1), i.e. with the second derivative of the time-averaged MSD. Hence

$$C(\tau) \propto \frac{d^2 \langle \mathbf{r}^2(\tau) \rangle_t}{d\tau^2} \propto \tau^{\alpha - 2} \; .$$

Despite the lack of full mathematical rigor, this heuristic argument suggests a general validity on the power-law decay observed in the ACVF, irrespective of being a FBM. However, it does not provide any hint on the range  $\xi \leq 1$  and hence also cannot claim anything on the integral area below the ACVF curve, i.e. these might depend considerably on the random walk process. But at least for the static percolation problem, all FBM features of the ACVF appear to be met.





Universität Bayreuth • 95440 Bayreuth

PCCP

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence

Open Access Article. Published on 17 June 2025. Downloaded on 6/18/2025 8:44:35 AM.

**Editorial Office** 

View Article Online LEHRSTUHL FÜROOI: 10.1039/D5CP01378J EXPERIMENTALPHYSIK I

Prof. Dr. Matthias Weiss

Universität Bayreuth Universitätsstraße 30 95447 Bayreuth

Telefon: +49 (0) 921 / 55 - 2500 Internet: http://www.ep1.uni-bayreuth.de e-mail: matthias.weiss@uni-bayreuth.de

The submitted manuscript is based on simulation data.

Data analyses and trajectory calculation for fractional Brownian motion were done with a previously published toolbox of Matlab codes, see Rehfeldt & Weiss, Soft Matter **19**, 5206 (2023) and the associated GitHub repository https://github.com/mweisslab/sptanalysis.

Fortran90 codes that were used to obtain data for obstructed diffusion are also included and available in this GitHub repository as of April 10, 2025.